arXiv:1407.2957v1 [math.NT] 9 Jul 2014

A NOTE ON BOOLE POLYNOMIALS WITH q-PARAMETER

DAE SAN KIM, YU SEON JANG, TAEKYUN KIM, AND SEOG-HOON RIM,

Abstract. Recently, Boole polynomials have been studied by Kim and Kim over the p-adic number field. In this paper, we consider a q-extension of Boole polynomials by using the fermionic p-adic integrals on $\mathbb{Z}_p$ and give some new identities related to those polynomials.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will denote the ring of p-adic integers, the field of p-adic numbers and the completion of algebraic closure of $\mathbb{Q}_p$. The p-adic norm $|\cdot|_p$ is normalized as $|p|_p = 1/p$. Let $C(\mathbb{Z}_p)$ be the space of continuous functions on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, the fermionic p-adic integral on $\mathbb{Z}_p$ is defined by Kim to be

$$I(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)(-1)^x, \text{ (see [11])}. \tag{1}$$

From (1), we have

$$I_{-1}(f_n) = 2\sum_{x=0}^{n-1}(-1)^{n-1-a} f(a) + (-1)^n I_{-1}(f). \tag{2}$$

Let us assume that q is an indeterminate in $\mathbb{C}_p$ with $|1-q|_p < p^{-1/(p-1)}$. The Stirling number of the first kind is defined by

$$(x)_n = \sum_{\ell=0}^{n} S_1(n,\ell) x^\ell, \ (n \geq 0), \tag{3}$$

and the Stirling number of the second kind is given by

$$x^n = \sum_{\ell=0}^{n} S_2(n,\ell)(x)_\ell, \ (n \geq 0), \text{ (see [9, 16])}. \tag{4}$$

The Boole polynomials are defined by the generating function to be

$$\sum_{n=0}^{\infty} Bl_n(x|\lambda)\frac{t^n}{n!} = \frac{1}{1+(1+t)^{\lambda}}(1+t)^x, \text{ (see [9, 16])}. \tag{5}$$

In [9], Kim and Kim gave a Witt-type formula for $Bl_n(x|\lambda)$ over the p-adic number field as follows:

$$\int_{\mathbb{Z}_p} (x+\lambda y)_n d\mu_{-1}(y) = 2Bl_n(x|\lambda), \ (n \geq 0), \tag{6}$$

where $\lambda \in \mathbb{Z}_p$ and $(x)_n = x(x-1)\cdots(x-n+1)$.

Let us define the q-product of x as follows:

$$(x)_{n,q} = x(x-q)(x-2q)\cdots(x-(n-1)q), \ (n \geq 0). \tag{7}$$

As is known, the Euler polynomials are defined by

$$\frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}, \text{ (see [1 - 20])}. \tag{8}$$

When $x = 0, E_n = E_n(0)$ are called the Euler numbers. From (2), we have

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) = \frac{2}{e^t+1}e^{xt} = \sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!}. \tag{9}$$

In this paper, we consider a q-extension of Boole polynomials by using the fermionic p-adic integrals on $\mathbb{Z}_p$ and give some new identities of those polynomials.

2. Boole polynomials with q-parameter

In this section, we assume that $t \in \mathbb{C}_p$ with $|t|_p < p^{-1/(p-1)}|q|_p$ and $\lambda \in \mathbb{Z}_p$. Now, we consider the Boole polynomials with q-parameter as follows:

$$Bl_{n,q}(x|\lambda) = \int_{\mathbb{Z}_p} (x+\lambda y)_{n,q}\, d\mu_{-1}(y), \ (n \geq 0). \tag{10}$$

Thus, by (10), we get

$$\begin{aligned} Bl_{n,q}(x|\lambda) &= \sum_{\ell=0}^{n} S_1(n,\ell) q^{n-\ell}\lambda^{\ell} \int_{\mathbb{Z}_p} \left(\frac{x}{\lambda}+y\right)^{\ell} d\mu_{-1}(y) \\ &= \sum_{\ell=0}^{n} S_1(n,\ell) q^{n-\ell}\lambda^{\ell} E_{\ell}\left(\frac{x}{\lambda}\right). \end{aligned} \tag{11}$$

From (2) and (10), we can derive the generating function of $Bl_{n,q}(x|\lambda)$ as follows:

$$\begin{aligned}\sum_{n=0}^{\infty} Bl_{n,q}(x|\lambda)\frac{t^n}{n!} &= \sum_{n=0}^{\infty}\int_{\mathbb{Z}_p}(x+\lambda y)_{n,q}d\mu_{-1}(y)\frac{t^n}{n!}\\ &= \sum_{n=0}^{\infty} q^n \int_{\mathbb{Z}_p}\binom{\frac{x+\lambda y}{q}}{n}d\mu_{-1}(y)t^n\\ &= \int_{\mathbb{Z}_p}(1+qt)^{\frac{x+\lambda y}{q}}d\mu_{-1}(y)\\ &= (1+qt)^{\frac{x}{q}}\left(\frac{2}{(1+qt)^{\lambda/q}+1}\right).\end{aligned} \tag{12}$$

Therefore, by (12), we obtain the following theorem.

Theorem 2.1. *Let $F(t,x|\lambda)=\sum_{n=0}^{\infty} Bl_{n,q}(x|\lambda)\frac{t^n}{n!}$. Then we have*

$$F(t,x|\lambda)=\left(\frac{2}{(1+qt)^{\lambda/q}+1}\right)(1+qt)^{x/q}.$$

By replacing t by $(e^t-1)/q$ in (12), we get

$$\begin{aligned}\sum_{n=0}^{\infty} Bl_{n,q}(x|\lambda)q^{-n}\frac{(e^t-1)^n}{n!} &= \frac{2}{e^{\lambda t/q}+1}e^{\frac{x}{q}t}\\ &= \sum_{n=0}^{\infty} E_n\left(\frac{x}{\lambda}\right)\left(\frac{\lambda}{q}\right)^n\frac{t^n}{n!}\end{aligned} \tag{13}$$

and

$$\begin{aligned}\sum_{n=0}^{\infty} Bl_{n,q}(x|\lambda)q^{-n}\frac{(e^t-1)^n}{n!} &= \sum_{n=0}^{\infty} Bl_{n,q}(x|\lambda)q^{-n}\sum_{m=n}^{\infty} S_2(m,n)\frac{t^m}{m!}\\ &= \sum_{m=0}^{\infty}\left\{\sum_{n=0}^{m} Bl_{n,q}(x|\lambda)\frac{S_2(m,n)}{q^n}\right\}\frac{t^m}{m!}.\end{aligned} \tag{14}$$

Therefore, by (13) and (14), we obtain the following theorem.

Theorem 2.2. *For $m\geq 0$, we have*

$$\lambda^m E_m\left(\frac{x}{\lambda}\right)=\sum_{n=0}^{m} Bl_{n,q}(x|\lambda)q^{m-n}S_2(m,n),$$

and

$$Bl_{m,q}(x|\lambda)=\sum_{\ell=0}^{m} S_1(m,\ell)q^{m-\ell}\lambda^{\ell}E_{\ell}\left(\frac{x}{\lambda}\right).$$

Note that $\lim_{q\to 1} Bl_{n,q}(x|\lambda)=2Bl_n(x|\lambda)$, $(n\geq 0)$. When $x=0, Bl_{n,q}(\lambda)=Bl_{n,q}(0|\lambda)$ are called the q-Boole numbers.

Now, we consider the q-Boole polynomials of the second kind as follows:

$$\widehat{Bl}_{n,q}(x|\lambda) = \int_{\mathbb{Z}_p} (-\lambda y + x)_{n,q} d\mu_{-1}(y), \ (n \geq 0). \tag{15}$$

Thus, by (15), we get

$$\begin{aligned} \widehat{Bl}_{n,q}(x|\lambda) =& q^n \int_{\mathbb{Z}_p} \left(\frac{-\lambda y + x}{q} \right)_n d\mu_{-1}(y) \\ =& q^n \int_{\mathbb{Z}_p} \sum_{\ell=0}^{n} \frac{\lambda^\ell S_1(n,\ell)}{q^\ell} (-1)^\ell \left(y - \frac{x}{\lambda} \right)^\ell d\mu_{-1}(y) \\ =& \sum_{\ell=0}^{n} S_1(n,\ell) q^{n-\ell} \lambda^\ell (-1)^\ell E_\ell \left(-\frac{x}{\lambda} \right). \end{aligned} \tag{16}$$

From (8), we have

$$\begin{aligned} \sum_{n=0}^{\infty} E_n \left(-\frac{x}{\lambda} \right) \frac{t^n}{n!} =& \frac{2}{e^t + 1} e^{(-\frac{x}{\lambda})t} \\ =& \frac{2}{1 + e^{-t}} e^{-\left(1 + \frac{x}{\lambda}\right)t} \\ =& \sum_{n=0}^{\infty} (-1)^n E_n \left(1 + \frac{x}{\lambda} \right) \frac{t^n}{n!}. \end{aligned} \tag{17}$$

By (16) and (17), we get

$$\widehat{Bl}_{n,q}(x|\lambda) = \sum_{\ell=0}^{n} \lambda^\ell |S_1(n,\ell)| q^{n-\ell} E_\ell \left(1 + \frac{x}{\lambda} \right). \tag{18}$$

From (15), we can derive the generating function of the Boole polynomials of the second kind as follows:

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{Bl}_{n,q}(x|\lambda) \frac{t^n}{n!} =& \sum_{n=0}^{\infty} q^n \int_{\mathbb{Z}_p} \binom{\frac{-\lambda y + x}{q}}{n} d\mu_{-1}(y) t^n \\ =& \int_{\mathbb{Z}_p} (1 + qt)^{\frac{-\lambda y + x}{q}} d\mu_{-1}(y) \\ =& (1 + qt)^{\frac{x}{q}} \int_{\mathbb{Z}_p} (1 + qt)^{-\frac{\lambda y}{q}} d\mu_{-1}(y) \\ =& (1 + qt)^{\frac{x+\lambda}{q}} \frac{2}{(1 + qt)^{\lambda/q} + 1}. \end{aligned} \tag{19}$$

By replacing t by $(e^t-1)/q$ in (19), we get

$$\begin{aligned}\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}(x|\lambda)\frac{1}{q^n}\frac{(e^t-1)^n}{n!} &= e^{\frac{1}{q}(x+\lambda)t}\frac{2}{e^{\frac{\lambda}{q}t}+1}\\ &= \sum_{n=0}^{\infty}E_n\left(\frac{x+\lambda}{\lambda}\right)\frac{\lambda^n}{q^n}\frac{t^n}{n!}\end{aligned} \tag{20}$$

and

$$\begin{aligned}\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}(x|\lambda)q^{-n}\frac{(e^t-1)^n}{n!} &= \sum_{n=0}^{\infty}\widehat{Bl}_{n,q}(x|\lambda)q^{-n}\sum_{m=n}^{\infty}S_2(m,n)\frac{t^m}{m!}\\ &= \sum_{m=0}^{\infty}\left(\sum_{n=0}^{m}\widehat{Bl}_{n,q}(x|\lambda)\frac{S_2(m,n)}{q^n}\right)\frac{t^m}{m!}.\end{aligned} \tag{21}$$

Therefore, by (18), (20) and (21), we obtain the following theorem.

Theorem 2.3. *For $m \geq 0$, we have*

$$\sum_{n=0}^{m}\widehat{Bl}_{n,q}(x|\lambda)q^{m-n}S_2(m,n) = \lambda^m E_m\left(1+\frac{x}{\lambda}\right)$$

and

$$\widehat{Bl}_{m,q}(x|\lambda) = \sum_{\ell=0}^{m}S_1(m,\ell)q^{m-\ell}\lambda^{\ell}E_{\ell}\left(1+\frac{x}{\lambda}\right).$$

For $\alpha \in \mathbb{N}$, let us consider q-Boole polynomials of the first kind with order α as follows:

(22)

$$\begin{aligned}Bl_{n,q}^{(\alpha)}(x|\lambda) &= \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(\lambda x_1+\cdots+\lambda x_\alpha+x)_{n,q}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha)\\ &= q^n\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}\left(\frac{\lambda x_1+\cdots+\lambda x_\alpha+x}{q}\right)_n d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha)\\ &= q^n\sum_{\ell=0}^{n}S_1(n,\ell)\frac{1}{q^\ell}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(\lambda x_1+\cdots+\lambda x_\alpha+x)^{\ell}\,d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha)\\ &= \sum_{\ell=0}^{n}S_1(n,\ell)q^{n-\ell}\lambda^{\ell}E_{\ell}^{(\alpha)}\left(\frac{x}{\lambda}\right),\end{aligned}$$

where $E_n^{(\alpha)}(x)$ are the Euler polynomials of order α which are defined by

$$\left(\frac{2}{e^t+1}\right)^{\alpha}e^{xt} = \sum_{n=0}^{\infty}E_n^{(\alpha)}(x)\frac{t^n}{n!}.$$

From (22), we note that the generating function of $Bl_{n,q}^{(\alpha)}(x|\lambda)$ are given by

$$\begin{aligned}\sum_{n=0}^{\infty} Bl_{n,q}^{(\alpha)}(x|\lambda)\frac{t^n}{n!} &= \sum_{n=0}^{\infty} q^n \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}\binom{\frac{\lambda x_1+\cdots+\lambda x_\alpha+x}{q}}{n} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha)t^n \\ &= \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+qt)^{\frac{\lambda x_1+\cdots+\lambda x_\alpha+x}{q}} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha) \\ &= (1+qt)^{\frac{x}{q}}\left(\frac{2}{(1+qt)^{\frac{\lambda}{q}}+1}\right)^{\alpha}.\end{aligned} \tag{23}$$

By replacing t by $(e^t-1)/q$, we get

$$\begin{aligned}\sum_{n=0}^{\infty} Bl_{n,q}^{(\alpha)}(x|\lambda)\frac{1}{q^n}\frac{(e^t-1)^n}{n!} &= e^{\frac{x}{q}t}\left(\frac{2}{e^{\frac{\lambda}{q}t}+1}\right)^{\alpha} \\ &= \sum_{n=0}^{\infty} E_n^{(\alpha)}\left(\frac{x}{\lambda}\right)\frac{\lambda^n}{q^n}t^n\end{aligned} \tag{24}$$

and

$$\begin{aligned}\sum_{n=0}^{\infty} Bl_{n,q}^{(\alpha)}(x|\lambda)\frac{1}{q^n}\frac{(e^t-1)^n}{n!} &= \sum_{n=0}^{\infty} Bl_{n,q}^{(\alpha)}(x|\lambda)\frac{1}{q^n}\sum_{m=n}^{\infty} S_2(m,n)\frac{t^m}{m!} \\ &= \sum_{m=0}^{\infty}\left(\sum_{n=0}^{m} Bl_{n,q}^{(\alpha)}(x|\lambda)\frac{S_2(m,n)}{q^n}\right)\frac{t^m}{m!}.\end{aligned} \tag{25}$$

Therefore, by (24) and (25), we obtain the following theorem.

Theorem 2.4. *For $m \geq 0$, we have*

$$\lambda^m E_m^{(\alpha)}\left(\frac{x}{\lambda}\right) = \sum_{n=0}^{m} q^{m-n} Bl_{n,q}^{(\alpha)}(x|\lambda)S_2(m,n)$$

and

$$Bl_{m,q}^{(\alpha)}(x|\lambda) = \sum_{\ell=0}^{m} S_1(m,\ell)q^{m-\ell}\lambda^\ell E_\ell^{(\alpha)}\left(\frac{x}{\lambda}\right).$$

We consider the q-Boole polynomials of the second kind with order α as follows:

$$\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda) = \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(-\lambda x_1-\cdots-\lambda x_\alpha+x)_{n,q}\, d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha). \tag{26}$$

Then, by (26), we get

$$\begin{aligned}\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda) &= q^n\sum_{\ell=0}^{n} S_1(n,\ell)(-1)^\ell\left(\frac{\lambda}{q}\right)^\ell\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}\left(x_1+\cdots+x_\alpha-\frac{x}{\lambda}\right)^\ell d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha) \\ &= \sum_{\ell=0}^{n} S_1(n,\ell)(-1)^\ell q^{n-\ell}\lambda^\ell E_\ell^{(\alpha)}\left(-\frac{x}{\lambda}\right).\end{aligned} \tag{27}$$

From the definition of the higher-order Euler polynomials, we note that

$$\begin{aligned}\sum_{n=0}^{\infty} E_n^{(\alpha)}\left(-\frac{x}{\lambda}\right)\frac{t^n}{n!} &= \left(\frac{2}{e^t+1}\right)^{\alpha} e^{-\frac{x}{\lambda}t} \\ &= \left(\frac{2}{1+e^{-t}}\right)^{\alpha} e^{-\left(\frac{x}{\lambda}+\alpha\right)t} \\ &= \sum_{n=0}^{\infty}(-1)^n E_n^{(\alpha)}\left(\frac{x}{\lambda}+\alpha\right)\frac{t^n}{n!}.\end{aligned} \tag{28}$$

Thus, by (27) and (28), we get

$$\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda) = \sum_{\ell=0}^{n} S_1(n,\ell) q^{n-\ell}\lambda^{\ell} E_{\ell}^{(\alpha)}\left(\frac{x}{\lambda}+\alpha\right). \tag{29}$$

From (26), we have

(30)

$$\begin{aligned}\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda)\frac{t^n}{n!} &= \sum_{n=0}^{\infty} q^n \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}\binom{\frac{-\lambda x_1+\cdots-\lambda x_\alpha+x}{q}}{n} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha)t^n \\ &= \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+qt)^{\frac{x-\lambda x_1-\cdots-\lambda x_\alpha}{q}} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_\alpha) \\ &= (1+qt)^{\frac{x+\alpha}{q}}\left(\frac{2}{(1+qt)^{\frac{\lambda}{q}}+1}\right)^{\alpha}.\end{aligned}$$

By replacing t by $(e^t-1)/q$, we get

$$\begin{aligned}\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda)\frac{(e^t-1)^n}{n!q^n} &= e^{\frac{x+\alpha}{q}t}\left(\frac{2}{e^{\frac{\lambda}{q}t}+1}\right)^{\alpha} \\ &= \sum_{n=0}^{\infty} E_n^{(\alpha)}\left(\frac{x+\alpha}{\lambda}\right)\left(\frac{\lambda}{q}\right)^n\frac{t^n}{n!}.\end{aligned} \tag{31}$$

and

$$\begin{aligned}\sum_{n=0}^{\infty}\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda)\frac{1}{q^n}\frac{1}{n!}(e^t-1)^n &= \sum_{n=0}^{\infty}\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda)\frac{1}{q^n}\sum_{m=n}^{\infty} S_2(m,n)\frac{t^m}{m!} \\ &= \sum_{m=0}^{\infty}\left(\sum_{n=0}^{m}\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda)\frac{S_2(m,n)}{q^n}\right)\frac{t^m}{m!}.\end{aligned} \tag{32}$$

Therefore, by (31) and (32), we obtain the following theorem.

Theorem 2.5. *For $m \geq 0$, we have*

$$\lambda^m E_m^{(\alpha)}\left(\frac{x+\alpha}{\lambda}\right) = \sum_{n=0}^{m} q^{m-n}\widehat{Bl}_{n,q}^{(\alpha)}(x|\lambda) S_2(m,n) \tag{33}$$

and

$$\widehat{Bl}_{m,q}^{(\alpha)}(x|\lambda)=\sum_{\ell=0}^{m}S_1(m,\ell)q^{m-\ell}\lambda^{\ell}E_{\ell}^{(\alpha)}\left(\frac{x+\alpha}{\lambda}\right). \tag{34}$$

Remark. When $x=0$, $\widehat{Bl}_{n,q}(\lambda)=\widehat{Bl}_{n,q}(0|\lambda)$ are called the q-Boole numbers of the second kind.

Now, we observe that

$$\begin{aligned}\frac{\widehat{Bl}_{n,q}(\lambda)}{n!}=&\frac{1}{n!}\int_{\mathbb{Z}_p}(-\lambda y)_{n,q}d\mu_{-1}(y)\\ =&q^n\int_{\mathbb{Z}_p}\binom{\frac{-\lambda y}{q}}{n}d\mu_{-1}(y)\\ =&(-1)^nq^n\int_{\mathbb{Z}_p}\binom{\frac{\lambda y}{q}+n-1}{n}d\mu_{-1}(y)\\ =&(-1)^nq^n\sum_{\ell=0}^{n}\binom{n-1}{\ell-1}\frac{1}{\ell!}\int_{\mathbb{Z}_p}\left(\frac{\lambda y}{q}\right)_{\ell}d\mu_{-1}(y)\\ =&(-1)^nq^n\sum_{\ell=0}^{n}\binom{n-1}{\ell-1}\frac{Bl_{\ell,q}(\lambda)}{\ell!q^{\ell}}.\end{aligned} \tag{35}$$

Therefore, by (35), we obtain the following theorem.

Theorem 2.6. *For $n\geq 0$, we have*

$$\frac{(-1)^n}{q^n}\frac{\widehat{Bl}_{n,q}(\lambda)}{n!}=\sum_{\ell=0}^{n}\binom{n-1}{\ell-1}\frac{Bl_{\ell,q}(\lambda)}{\ell!q^{\ell}}. \tag{36}$$

References

[1] S. Araci and M. Acikgoz, A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials, *Adv. Stud. Contemp. Math.* **22** (2012), no. 3, 399–406.

[2] A. Bayad and J. Chikhi, Apostol-Euler polynomials and asymptotics for negative binomial reciprocals, *Adv. Stud. Contemp. Math.* **24** (2014), no. 1, 33–37.

[3] M. Can, M. Cenkci, V. Kurt, and Y. Simsek, Twisted Dedekind type sums associated with Barnes' type multiple Frobenius-Euler ℓ-functions, *Adv. Stud. Contemp. Math.* **18** (2009), no. 2, 135–160.

[4] J. Choi, D. S. Kim, T. Kim, and Y. H. Kim, Some arithmetic identities on Bernoulli and Euler numbers arising from the p-adic integrals on $\mathbb{Z}_p$, *Adv. Stud. Contemp. Math.* **22** (2012), no. 2, 239–247.

[5] D. Ding and J. Yang, Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials, *Adv. Stud. Contemp. Math.* **20** (2010), no. 1, 7–21.

[6] D. V. Dolgy, T. Kim, B. Lee, and C. S. Ryoo, On the q-analogue of Euler measure with weight α, *Adv. Stud. Contemp. Math.* **21** (2011), no. 4, 429–435.

[7] S. Gaboury, R. Tremblay, and B.-J. Fugère, Some explicit foumulas for certain new classes of Bernoulli, Euler and Genocchi polynomials, *Proc. Jangjeon Math. Soc.* **17** (2014), no. 1, 115–123.

[8] J. H. Jeong, J.-H. Jin, J.-W. Park, and S.-H. Rim, On the twisted weak q-Euler numbers and polynomials with weight 0, *Proc. Jangjeon Math. Soc.* **16** (2013), no. 2, 157–163.
[9] D. S. Kim and T. Kim, A note on Boole polynomials, *Integral Transforms Spec. Funct.* **25** (2014), no. 8, 627–633.
[10] T. Kim, D. S. Kim, D. V. Dolgy, and S.-H. Rim, Some identities on the Euler numbers arising from Euler basis polynomials, *Ars Combin.* **109** (2013), 433–446.
[11] T. Kim, Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$, *Russ. J. Math. Phys.* **16** (2009), no. 1, 93–96.
[12] T. Kim, Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$, *Russ. J. Math. Phys.* **16** (2009), no. 4, 484–491.
[13] T. Kim, A study on the q-Euler numbers and the fermionic q-integral of the product of several type q-Bernstein polynomials on $\mathbb{Z}_p$, *Adv. Stud. Contemp. Math.* **23** (2013), no. 1, 5–11.
[14] Q.-M. Luo, q-analogues of some results for the Apostol-Euler polynomials, *Adv. Stud. Contemp. Math.* **20** (2010), no. 1, 103–113.
[15] H. Ozden, I. N. Cangul, and Y. Simsek, Remarks on q-Bernoulli numbers associated with Daehee numbers, *Adv. Stud. Contemp. Math.* **18** (2009), no. 1, 41–48.
[16] S. Roman, The umbral calculus, Academic Press, Inc., New York, 1984.
[17] C. S. Ryoo, H. Song and R. P. Agarwal, On the roots of the q-analogue of Euler-Barnes' polynomials, *Adv. Stud. Contemp. Math.* **9** (2004), no. 2, 153–163.
[18] Y. Simsek, O. Yurekli and V. Kurt, On interpolation functions of the twisted generalized Frobenius-Euler numbers, *Adv. Stud. Contemp. Math.* **15** (2007), no. 2, 187–194.
[19] E. Sen, Theorems on Apostol-Euler polynomials of higher order arising from Euler basis, *Adv. Stud. Contemp. Math.* **23** (2013), no. 2, 337–345.
[20] Z. Zhang and H. Yang, Some closed formulas for generalized Bernoulli-Euler numbers and polynomials, *Proc. Jangjeon Math. Soc.* **11** (2008), no. 2, 191–198.

D. S. Kim
Department of Mathematics
Sogang Uiversity
Seoul, Republic of Korea
e-mail: dskim@sogang.ac.kr

Y. S. Jang
Department of Applied Mathematics
Kangnam University
Yongin 446-702, Republic of Korea
e-mail: ysjang@kangnam.ac.kr

T. Kim
Department of Mathematics
Kwangwoon University
Seoul 139-701, Republic of Korea
e-mail: tkkim@kw.ac.kr

S.-H. Rim
Department of Mathematics Education

Kyungpook National University
Taegu 702-701, Republic of Korea
e-mail: shrim@knu.ac.kr